

AGPRO

RAINBOW

WARNING
Keep out of reach of children

RYDAZIN

HERBICIDE

Selective pre-and post-emergent herbicide for fodder, beet, silver beet, red beet, sugar beet and mangolds, and post emergent weed control in onions and chives.

Contains 430 g/L chloridazon in the form of a suspension concentrate

Registered pursuant to the ACVM Act 1997 No:P9353
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to HSNO Act 1996 No HSR101123
See www.epa.govt.nz for approval conditions

Distributed by:
AGPRO NZ LTD
10 Polaris Place, East Tamaki, Auckland
PO Box 58 963, Botany
Phone: 09 273 3456
Email: sales@agpro.co.nz
Website: www.agpro.co.nz

Net contents: 10L

Batch number:

HAZARD CLASSIFICATIONS:

Classifications: 6.5B, 9.1A, 9.2A

HEALTH WARNING: May be harmful if swallowed, inhaled or absorbed through the skin. May cause sensitization from prolonged skin contact. Avoid skin contact and inhalation of spray mist.

ENVIRONMENTAL WARNING

VERY TOXIC TO AQUATIC ORGANISMS. Avoid contamination of any water supply with product or empty container. **VERY TOXIC TO THE SOIL ENVIRONMENT:** Herbicide.

HEALTH WARNING: HARMFUL. May be harmful if swallowed, inhaled or absorbed through the skin. May cause sensitization from prolonged skin contact. Avoid skin contact and inhalation of spray mist.

DISPOSAL: Container: Triple rinse empty container and add residue to the spray tank. Recycle through an approved recycling facility. Product: Dispose of this product only by using according to the label or at an approved landfill. Do NOT burn product. Do NOT contaminate any water supply with product or used container.

HANDLING PRECAUTIONS: When mixing or applying product, wear protective clothing, glove and goggles. Do not eat or drink while using. Wash hands and face before meals and after work. Wash protective clothing after work. Store in original container away from food and feedstuffs. Flush spray equipment thoroughly after use.

APPROVED HANDLER: This product must be under the control of an approved handler if applied in a widely dispersed manner or if applied by a contractor.

STORAGE: Store in original container, tightly closed, in a locked, dry, cool, well-ventilated place away from feed, seeds and foodstuffs. Storage must be generally in accordance with NZS 8904 Management of Agrichemicals. See the HAZNOTE for further information. When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

First Aid

IF SWALLOWED: Do NOT induce vomiting. Rinse mouth. For advice contact the National Poisons Centre 0800 POISONS (0800 764 766) or a doctor. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water.

PRODUCT INFORMATION

RYDAZIN is a selective pre- and post-emergence herbicide providing a good residual action against a wide range of annual broadleaf weeds and some annual grasses with a contact action on a wide range of weeds up to the 2 leaf stage. Adequate soil moisture is required for good residual activity. The post-emergent effect can be enhanced with the addition of paraffinic crop oil spraying oil, RYDAZIN remains effective in the soil for several weeks.

Susceptible weeds		Moderately Susceptible	Resistant
Annual poa	Penny cress	Barnyard grass	Couch
Black nightshade	Rayless chamomile	Cleavers	Thistles
Charlock	Redroot	Curled dock	All weeds past
Chick weed	Speedwells	Fumitory	the 2 leaf stage
Docks	Water purslane	Spurrey	
Fathen	Wild carrot	Summer grass	
Galinsoga	Wild radish		
Groundsel	Willow weed		

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

DIRECTIONS FOR USE

Mixing: Always shake the container before use. Pour the required amount directly into the partly filled spray tank with agitation running. Complete filling of the spray tank.

Recommendations: Apply RYDAZIN in 300-600 liters of water per hectare. Use the higher rate on heavy clay soils or soils rich in organic matter.

FODDER BEET, RED BEET, SILVER BEET, MANGOLDS:

PRE-EMERGENCE: Apply 7 to 10.5 litres per hectare as soon as possible after seeding and before crop emergence, to a fine, moist soil surface. If weeds are present, mix with a suitable contact herbicide. If soil is dry, rain or irrigation is required within 6 to 7 days for residual activity.

POST-EMERGENCE: Apply 7-9 litres per hectare to a weed free soil after the first true leaf is visible on the beet. Do not spray in the 8 days before thinning. The addition of paraffinic crop oil spraying oil at 1 litre per hectare will enhance the knockdown on weed up to 2 true leaf.

ONIONS AND CHIVES:

POST-EMERGENCE: Apply 1 to 2 litres RYDAZIN with 1 to 2 litres of Chlorpropham (400g/L formulation) and 1 litre of paraffinic crop oil spraying oil per hectare at the flag stage. Apply the RYDAZIN and the chlorpropham at equal rates for best results. Repeat the application 2 to 3 weeks later. Weeds should not have passed the 4 true leaf stage. For best contact action, no rain should fall for 6 hours after application. Rain is required, however, within 7 days for good residual activity.

CAUTIONS: Crop damage may occur on light soils.

Spray Drift: Beware- apply this product carefully. Spray drift may cause serious damage to desirable plants. Ensure compliance with any relevant regional air management plants.

Registered to:

Shandong Rainbow International Co., Ltd
6th Floor, Gaoxinjishuchuangye Centre
Shunhua Road 750#, Jinan
Shandong, 250101, China

NZ Agent:

AR & JA DRYSDALE LTD
PO Box 72275
Papakura 2244
Tel: (021) 939 728

CONDITIONS OF SALE: The use of RYDAZIN herbicide being beyond the control of the suppliers, no warranty expressed or implied, beyond that covered under the Fair Trading Act is given by the supplier regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and the supplier accepts no responsibility for any consequences whatsoever resulting from the use of this product.