

WARNING

KEEP OUT OF REACH OF CHILDREN

Read label before use

ECOTOXIC

Afupet

***Active Ingredient: contains 500 g/L Folpet
in the form of a suspension concentrate.***

For the control of speckled leaf blotch (*Septoria tritici*) in
wheat and scald and Ramularia in barley

10L

Registered pursuant to the ACVM Act 1997, No. P10181
See www.foodsafety.govt.nz for registration conditions
Approved pursuant to HSNO Act 1996,
Approval Code HSR101068
See www.epa.govt.nz for approval controls

Distributed by:
AGPRO (N.Z.) Ltd
10 Polaris Place, East Tamaki, Auckland 2013
PO Box 58-963, Botany, Auckland 2163
Ph 09 273 3456
www.agpro.co.nz

LBL100.1F

AGPRO



HAZARD CLASSIFICATIONS:

Carcinogenicity Category 2,
Hazardous to terrestrial vertebrates,
Hazardous to the aquatic environment
acute Category 1

TOXICITY

WARNING - Suspected of causing
cancer.

ECOTOXICITY

Very toxic to aquatic organisms.
Harmful to terrestrial vertebrates.
Do not apply directly onto or into
water. Take all reasonable steps to
ensure that the substance does not
cause any significant adverse effects
to the environment beyond the
application area.

QUALIFIED HANDLER: Anyone who
mixes, loads, or applies this product
is required to be suitably qualified,
or must receive guidance and, if
required, assistance from a suitably
qualified person.

RECORD KEEPING:

Where 3 kg or more of **Afupet** is
applied within 24 hours, in a place
where the substance is likely to enter
air or water and leave the application
area, a written record is required to
be kept of each application.

Batch Number:

Date of Manufacture:

PRECAUTIONS

Keep out of reach of children. Obtain
special instructions before use.

Do not handle until all safety
precautions have been read and
understood. Use personal protective
equipment as required.

PERSONAL PROTECTION

When mixing or applying wear
appropriate protective clothing
including cotton overalls buttoned to
the neck and wrist, impervious, elbow-
length gloves, and eye protection.
Remove protective clothing and wash
hands, arms and face with soap and
water before meals and after work.

Do not eat, drink or smoke when
using.

Wash protective clothing and
equipment before reuse.

STORAGE

Store locked up in the original,
unopened container in a cool, dry
place, out of direct sunlight and away
from stockfeed or foodstuffs.

Store in accordance with the current
version of NZS 8409 Management
of Agrichemicals. Secondary
containment, signage and an
emergency response plan are required
where 100 L or more of this product
are stored.



AGPRO[®]
(N.Z.) LIMITED



LBL100.1B

SHELF LIFE

When stored appropriately this product should show no significant degradation for two years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

DISPOSAL

Avoid contamination of any water supply with product or empty container.

Container Disposal - Triple rinse empty container and add rinsate to spray tank. Submit clean empty container to an agricultural recycling depot for recycling. Alternatively, puncture and bury in a suitable landfill.

Product Disposal - Dispose of this product only by using according to the label, or through an approved agricultural chemical disposal facility.

SPILLAGE

Collect spill. Exclude all bystanders from the vicinity of the spill and prevent material from entering waterways. Wear appropriate protective clothing as described in the personal protection section of this label. Absorb spills with an inert material (such as vermiculite, earth, sand or synthetic absorbent substance) and place in waste containers. Wash area with water and alkaline detergent. Then absorb any remaining liquid with further inert material. Dispose of the waste safely in an approved landfill.

EMERGENCY RESPONSE

In a transport emergency call 111, Police or Fire Brigade.

For specialist advice in a chemical emergency call 0800 CHEMCALL (0800 243 622) 24 hours.

SAFETY DATA SHEETS

Additional information is provided in the safety data sheet. Phone 09 273 3456 or visit www.agpro.co.nz for a copy.

FIRST AID

For advice, or if you are exposed or concerned, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If medical advice is needed, have product container or label at hand.

IF SWALLOWED: Do NOT induce vomiting. Rinse mouth. Get medical advice.

IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing. Get medical advice if you feel unwell.

IF ON SKIN: Wash with plenty of water. If skin irritation occurs get medical advice.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTRE or doctor for advice.

EQUIPMENT

Use accurately calibrated and maintained equipment in accordance with the New Zealand Standard for the Management of Agrichemicals (NZS8409). Flush equipment thoroughly with clean water after use. Do not use equipment to apply other pesticides to susceptible crops and plants. Dispose of cleaning water onto ground away from waterways.

PRODUCT INFORMATION

Afupet contains folpet, a protectant fungicide from the phthalimide chemical group with multi-site activity. For best results, thorough coverage of the crop prior to the onset of infection is recommended as **Afupet** only has

activity against very early stages of infection such as spore germination. Always tank-mix **Afupet** with an alternative mode of action fungicide such as Epoxiconazole 50G/L + Prochloraz 225G/L EC.

DIRECTIONS FOR USE

This product must not be applied at rates exceeding 1.5 L/ha and must not be applied in the same area more than two times in any 365 day period.

A minimum application interval of at least 14 days must be observed.

This product must be applied by ground-based application methods only. Do NOT use aerial application methods.

Do not apply this substance within 2 metres of a downwind water body. It is an offence to use this product on animals.

Mixing: Ensure concentrate is well-mixed before preparing spray solution. Shake or roll containers to mix contents. Add **Afupet** slowly to water in the spray tank with agitation. Continuous agitation is recommended to keep the product in suspension.

CROP	DISEASE	RATE	CRITICAL COMMENTS
Barley	Ramularia Scald	1.5 L/ha plus an alternative mode of action fungicide (e.g. Epoxiconazole 50g/L + Prochloraz 225g/L EC) in a minimum of 150 L of water/ha	Always apply Afupet in a tank-mixture with the full label rate of an alternative mode of action fungicide registered for the control of listed diseases. Apply up to two applications of Afupet per crop as part of a season-long fungicide spray program for disease management. The first application of Afupet should be as a preventative spray on susceptible cultivars and where conditions favour disease. Apply the first spray before disease becomes established in the crop. Re-apply in 3 - 4 weeks if disease pressure continues or if re-infection occurs. For scald control in barley, do not apply later than GS 39.
Wheat	Speckled leaf blotch		

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIOD

Barley: Do not apply after emergence of inflorescence completed (GS 59).

Wheat: Do not apply after flag leaf emergence (GS 39).

Grazing/stockfood: Do not graze or harvest barley and wheat crops for green feed (including silage) for 28 days after application.

COMPATIBILITY

Afupet is compatible with most insecticides and fungicides.

RESISTANCE MANAGEMENT

GROUP	M4	FUNGICIDE
--------------	-----------	------------------

Afupet contains the GROUP M4 phthalimide fungicide, folpet. Folpet is a multisite protectant fungicide at low risk of resistance development. However, unless directed, do not use another product containing a fungicide in the same group in conjunction with **Afupet**. Fungicides from different group codes should either be used alternately or mixed together to prevent the numbers of resistant individuals building up. To minimise this risk, use strictly in accordance with label instructions and resistance management strategies at www.resistance.nzpps.org/.

REGISTERED TO:

Shandong Rainbow International Co. Ltd,
30th Floor, Building A5-5,
Hanyu Financial Centre,
No. 7000 Jingshi Road,
Jinan Innovation Zone, Jinan,
Shandong, China

NZ Address:

8/8 Bagnall Ave., Point England,
Auckland 1072
Tel: +64 272664147

CONDITIONS OF SALE

As the use of this product is beyond the control of AGPRO (N.Z.) Ltd no warranty expressed or implied, beyond that covered under the Fair Trading Act is given by AGPRO (N.Z.) Ltd regarding its suitability, fitness, or efficacy for any purpose for which it is used by the user, whether in accordance with the directions or not. AGPRO (N.Z.) Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.