

WARNING

Keep out of reach of children

ECOTOXIC**AGPRO CHLORSULFURON 750WDG**

Contains: 750g/kg chlorsulfuron in the form of a water dispersible granule

For use in wheat, barley and oats to control most broadleaf weeds.

Read the directions for use before using this product.

Registered pursuant to the ACVM Act 1997: P008896
See www.foodsafety.govt.nz for registration conditions.
Approved pursuant to HSNO Act 1996 No HSR000231
See www.epa.govt.nz for approval conditions.

Date of manufacture:
Batch no:
Net contents: 500g

Registered to and distributed by:

AGPRO NZ Ltd
10 Polaris Place
East Tamaki
P O Box 58-963
East Tamaki
Auckland 2013
Ph 09 273 3456
Fax 09 273 3457
Website: www.agpro.co.nz

HAZARD CLASSIFICATION

6.3B, 9.1A, 9.1D, 9.2A

WARNING

ECOTOXIC: Harmful. May cause mild skin irritation. Very toxic to aquatic organisms and to some plant species.

PRECAUTIONS

APPROVED HANDLER: This product must be under the control of an approved handler when it is applied in a wide dispersive manner, or used by a commercial contractor. This product must not be used for any purpose or in any manner, contrary to this label unless authorised under appropriate legislation.

HANDLING PRECAUTIONS: Keep out of reach of children. Store in original container, tightly closed and in a locked, dry, cool area away from foodstuffs, fertilisers and seeds. Store in accordance with NZS 8409 Management of agrichemicals. Stores containing 100kg of AGPRO Chlorsulfuron 750WDG require signage and more than 100kg require emergency response plans.

PERSONAL PROTECTION: Avoid skin and eye contact and inhalation of spray mist. When mixing or applying wear appropriate eye protection, waterproof gloves, boots and overalls. Wash hands and exposed skin before meals and after work.

EQUIPMENT: Apply using well maintained and accurately calibrated equipment. Clean equipment thoroughly after use.

DISPOSAL: Triple rinse empty container and add rinsate to the spray tank. Recycle empty container. Otherwise crush and bury in a suitable landfill.

FIRST AID

If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair contact occurs, remove contaminated clothing and flush skin and hair with running water. If splashed in eyes, wash out immediately with water. If inhaled, move the victim to fresh air immediately.

SPILLAGE

Wear appropriate protective clothing and prevent material from entering waterways. Contain liquid spills and absorb with inert material. Sweep up solids without creating dust. Dispose of waste safely in a suitable landfill.

RECORD KEEPING

Records of use as described in NZS 8409 Management of Agrichemicals must be kept if more than 3kg of AGPRO Chlorsulfuron 750WDG is applied within 24 hours.

SHELF LIFE

When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

GENERAL INFORMATION

AGPRO Chlorsulfuron 750WDG is a selective herbicide for use in wheat, barley and oats to control most broadleaf weeds. It is non-corrosive, non-flammable and non-volatile. Best results are obtained when applied to young actively growing weeds. Symptoms (typically discoloration) may not be apparent for 1 – 3 weeks. Warm moist conditions promote the activity of AGPRO Chlorsulfuron 750WDG, whereas cold dry conditions delay the activity.

Weeds Controlled

Calandrinia, California thistle (Spray at the rosette stage (10 – 15 cm tall). Thistles that emerge after spraying will not be controlled.), chickweed, cornbind, dandelions, docks, fathen, hawksbeard, nodding thistle, rayless chamomile, redroot, scarlet pimpernel, scentless chamomile, scotch thistle, shepherds purse, spurrey, stinking mayweed, twin cress, vetch, white clover, willow weed, wild turnip, yellow gromwell.

Moderately Susceptible (Controlled if treated as seedlings under optimum conditions)

Cleavers, wireweed, yarrow.

Moderately Resistant (Weed vigour and density reduced)

Field pansy, speedwells.

RESISTANT WEEDS WARNING

Black nightshade, wild oats. To control tolerant weeds, tank mix a herbicide that is registered for control of these species. Follow label recommendations for the tank mix partner when mixing. AGPRO Chlorsulfuron 750WDG is a member of the sulfonyl urea group of herbicides. Naturally occurring biotypes resistant to sulfonyl urea herbicides are known to exist. These resistant weeds will not be controlled by AGPRO Chlorsulfuron 750WDG or other sulfonyl urea herbicides. To delay or prevent the development of resistant weeds, use AGPRO Chlorsulfuron 750WDG in tank mixes where appropriate, and or rotate with an effective herbicide of a different mode of action. AGPRO NZ Limited will not accept any liability for any losses that may result from use AGPRO Chlorsulfuron 750WDG failing to control resistant weeds, because it is difficult to determine the occurrence of weed resistance prior to use.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

DIRECTIONS FOR USE

Do not spray crops that are under stress from: drought, water saturated soil, disease or insect damage, as crop damage may occur. Crop damage may also occur if these conditions occur after spraying. Do not apply to crops undersown with lucerne or white clover.

Crop	Rate g/Ha	Critical Comments
Wheat, Barley, Oats	20	Apply after the crop is at the 2 leaf stage but before the boot stage. For best results, spray small (<5cm) actively growing weeds and before the crop canopy closes. A non-ionic surfactant must be added, use at label recommended rate.

MIXING INSTRUCTIONS

Using the graduated measuring cylinder provided, measure the correct amount of AGPRO Chlorsulfuron 750WDG required for the area to be sprayed and add to the required volume of water. Continuous agitation is required to keep the product in suspension. Use AGPRO Chlorsulfuron 750WDG spray mixtures within 24 hours of preparation, as product degradation may occur.

SURFACTANT

Always add a non-ionic surfactant at the manufacturer's recommendations unless mixing with another product which contains surfactant.

SPRAY VOLUME

Use a well maintained, accurately calibrated sprayer applying a minimum of 60 litres per hectare. Avoid overlapping and shut off spray booms while starting, slowing or stopping, or injury to the crop may result.

RE-CROPPING RECOMMENDATIONS

In the event of a crop failure, DO NOT replant to any other crop other than wheat or oats within 3 months of applying AGPRO Chlorsulfuron 750WDG. Following harvest of the cereal crop, and within 12 months of the AGPRO Chlorsulfuron 750WDG application, the paddock can be sown down into pasture, greenfeed crops (brassicas), peas or lucerne. DO NOT replant any other crop without first consulting AGPRO NZ Limited.

CONDITIONS OF SALE

The use of AGPRO Chlorsulfuron 750WDG herbicide being beyond the control of the manufacturer, no warranty expressed or implied, beyond that cover under the Fair Trading Act is given by AGPRO NZ Limited regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AGPRO NZ Limited accepts no responsibility for any consequence whatsoever resulting from the use of this product.