

**Keep Out Of Reach of Children  
Ecotoxic**

**NB: Read the directions and information on the label  
before using**



*[Signature]*  
16-11-2018

## **AGPRO Dimethomorph**

<b>Group</b>	<b>40</b>	<b>Fungicide</b>
--------------	-----------	------------------

Contains: 500 g/L dimethomorph in the form of a suspension concentrate.

AGPRO Dimethomorph is a fungicide for the control of Downy mildew in Grapes, Lettuce and Onions and Early and Late blight in Potatoes.

Registered pursuant to the ACVM Act 1997, No. P008705

See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

Approved pursuant to HSNO Act 1996, No. HSR007636

See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions.

Date of manufacture:

Batch Number:

Pack Size: 1 - 20 litres

Registered to and distributed by:

AGPRO NZ Ltd

10 Polaris Place

East Tamaki

P O Box 58-963

Auckland

Ph. 09 273 3456 Fax

09 273 3457

[www.agpro.co.nz](http://www.agpro.co.nz)

**HSNO Classifications: 6.9B, 9.1B**



*[Signature]*  
16-11-2018

**WARNING:**

Warning. May cause liver damage from repeated oral exposure at high doses. Toxic to the aquatic environment with long lasting effects.

**Record Keeping:** Records must be kept for 3 years.

**PRECAUTIONS:**

**Personal Protection**

Do not eat, drink or smoke whilst using.

Wear impermeable gloves, coat, hat and trousers during use. Use an air respirator with organic vapour canister or self contained breathing apparatus when exposure is likely.

Wear a face shield or goggles if eye contact is likely to occur.

Immediately after use or before any meal break, remove protective clothing and thoroughly wash hands and face. Avoid skin and eye contact.

**Storage**

Store in original container tightly closed and in a locked dry, cool, well ventilated area away from foodstuffs. Under appropriate storage, the product shows no significant degradation within 2 years from the date of manufacture. Contact the supplier if product is older than 2 years.

**Spills**

Wear the above mentioned protective clothing when dealing with spills. Contain the spill and recover spilt product with sawdust or other inert absorbent material then place recovered material into properly labelled drum. Deal with spillages immediately and prevent contamination of watercourses, drains, streams, ponds. If this is likely to occur, warn the local authorities immediately.

**Disposal**

Dispose of product by following label directions or at an approved landfill. Avoid contamination of waterways with product or empty container.

**Container Disposal**


Triple rinse container and add rinsate to spray tank. Burn rinsed container if circumstances, especially wind direction, are suitable. Otherwise bury in suitable landfill.

**FIRST AID:**

Contact National Poisons and Hazardous Chemicals Information Centre ph 0800 764 766 (0800 POISON) or a doctor for medical advice.

If swallowed, wash out mouth with water. Do not induce vomiting. Give a glass of water.

Transport patient to doctor or hospital as quickly as possible.



16-11-2018

If skin contact occurs, remove contaminated clothing and flush skin with soap and running water immediately. Wash contaminated clothing before re-use. If eye contact occurs, wash out eyes with running water immediately.

If inhaled, move person to fresh air immediately. Maintain half upright position if breathing is difficult. Carry out artificial respiration if necessary.

## GENERAL INFORMATION:

AGPRO Dimethomorph is a systemic Carboxylic acid amides (CAA) fungicide (group 40) with protectant, curative and anti-sporulant activity. AGPRO Dimethomorph should always be used as part of a protectant spray programme and be applied before disease symptoms appear.

## RESISTANCE MANAGEMENT

Resistance to AGPRO Dimethomorph and related CAA products could develop in some disease situations from repeated use of the product. To minimise this risk, use strictly in accordance with label instructions.

- Apply no more than four applications per crop.
- Under high disease pressure, apply 2 close interval applications of AGPRO Dimethomorph followed by applications of either protectant fungicides or fungicides with different modes of action.

**Application:** Apply using conventional spray equipment with thorough agitation. Complete spray coverage is essential for consistent disease control. The rates suggested are for high volume spraying to the point of run-off. Adjust dilution rate accordingly for concentrate spraying. Do not apply less than 200 L/ha water.

**Mixing:** Add the required quantity of AGPRO Dimethomorph to a half filled spray tank then fill to two thirds before adding surfactant. Complete filling and maintain agitation throughout filling and application.

**Compatibility:** Do NOT mix with strongly alkaline or acidic materials.

## Withholding Period:

Grapes, Lettuce, Onions, Potatoes                      14 days

## CONDITIONS OF SUPPLY

The use of AGPRO Dimethomorph being beyond the control of the manufacturer, no warranty expressed or implied, beyond that cover under the Fair Trading Act is given by AGPRO NZ Limited regarding suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AGPRO NZ Limited accepts no responsibility for any consequence whatsoever resulting from the use of this product.

**DIRECTIONS FOR USE:**

Crop	Disease	Rate	Remarks
Grapes	Downy mildew	36 mL/100 litres (360 mL/ha) + a suitable protectant fungicide at label rate	Apply AGPRO Dimethomorph in a minimum of 1000 litres water/ha. Apply at 14 day intervals from when shoots are 5 – 10 cm.
Lettuce	Downy mildew	360 mL/ha + a suitable protectant fungicide at label rate	Maintain a regular protectant programme. Apply AGPRO Dimethomorph at 7 to 10 day intervals when conditions favour disease development but before the disease is evident. Apply two consecutive sprays of AGPRO Dimethomorph 7 to 10 days apart and then change to a fungicide from another chemical group. Use the shorter interval when conditions favouring infection are creating a high risk of disease. Apply AGPRO Dimethomorph in 500-750 litres of water per hectare. Use a surfactant at labelled rates.
Onions	Downy mildew	360 mL/ha + a suitable protectant fungicide at label rate	Apply AGPRO Dimethomorph in 500-750 litres of water per hectare. Apply AGPRO Dimethomorph at 10 day intervals during periods of active growth or when high humidity conditions favour the disease. Use a surfactant at labelled rates.
Potatoes	Early blight Late blight	360 mL/ha + a suitable protectant fungicide at label rate	Apply AGPRO Dimethomorph as a foliar spray to run-off when the crop has fully emerged but before any sign of disease and repeat at 10-14 day intervals. Use the closer interval when conditions favour the disease, especially during favourable growth conditions in the period prior to flowering. Use a surfactant at labelled rates.



A handwritten signature in blue ink, appearing to be "Mark Brown", is written over the seal. Below the signature, the date "16-11-2018" is printed in a small font.

**It is an offence to use this product on animals.**

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**