WARNING

Keep Out of Reach of Children Read the label before use

AGPRO MOLLUS

All-weather Slug and Snail Bait

Contains 50 g/kg metaldehyde in the form of a bait.

Molluscicide for the control of Slugs and Snails in agricultural and horticultural situations

Pack size Kg

Registered pursuant to the ACVM Act 1997, No. P9735 See www.foodsafety.govt.nz for registration conditions Approved pursuant to HSNO Act 1996, Approval Code HSR007794 See www.epa.govt.nz for approval controls

> Registered to and distributed by: **AGPRO NZ Ltd** 10 Polaris Place, East Tamaki Auckland 2013 P O Box 58-963, Botany, Auckland 2163 Ph 09 273 3456 **www.agpro.co.nz**







HAZARD CLASSIFICATION Acute inhalation toxicity Category 4, Reproductive toxicity Category 2, Specific target organ toxicity (repeated exposure) Category 2, Hazardous to the aquatic environment chronic Category 3, Hazardous to terrestrial vertebrates.

HSNO Classification: 6.1D, 6.8B, 6.9B, 9.1C, 9.3C

This product must not be used for any purpose, or in any manner, contrary to this label unless authorised under appropriate legislation.

WARNING

TOXICITY

Harmful if inhaled. Suspected of damaging fertility or the unborn child. May cause damage to organs through prolonged or repeated exposure.

ECOTOXICITY

Harmful to aquatic life with long lasting effects. Harmful to terrestrial vertebrates. Do not apply directly into or onto water. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area. Avoid contamination of any water supply with product or empty packaging.

PRECAUTIONs

Obtain special instructions before use and do not handle until all safety precautions have been read and understood.

PERSONAL PROTECTION

Avoid contact with skin and eyes, ingestion and inhalation of dust.

When applying this product, overalls, boots, gloves, respiratory and eye protection should be worn. Do not eat, drink or smoke whilst using. Remove protective clothing and wash hands and face thoroughly before meals and after work.

STORAGE

Store in original container tightly closed and in a locked dry, cool, well-ventilated area avoiding direct sunlight and away from foodstuffs. Storage must be in accordance with the current version of NZ Standard for the Management of Agrichemicals (NZS 8409). Stores containing 1000 kg or more of **AGPRO Mollus** require signage and an emergency plan.



EQUIPMENT

Apply only with appropriate, well-maintained and calibrated equipment in accordance with NZS 8409.

SPILLAGE

Exclude non-essential people from the area. Wear protective clothing and prevent chemical from entering waterways. Generating as little dust as possible, sweep up solid spills and place in a sealable and labelled waste container. Wash area with water and absorb with inert material such as attapulgite, soil, etc., and place in the waste container. Dispose of safely at an approved landfill.

DISPOSAL

PRODUCT DISPOSAL: Dispose of this product only by using according to label directions or if this is not possible contact your supplier for advice.

PACKAGING DISPOSAL:

Dispose of empty packaging at an approved landfill. Do not burn. Do not use packaging for any other purpose.

FIRST AID

For advice call the National Poisons Centre 0800 POISON (0800 764 766) (24 hours) or a doctor. If medical advice is needed have product container or label to hand.

IF EXPOSED OR CONCERNED: Get medical advice/attention.

IF SWALLOWED: Do NOT induce vomiting. Rinse mouth. Call the Poisons Centre or a doctor if you feel unwell.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If irritation persists, get medical advice/attention.

IF ON SKIN OR HAIR: Remove contaminated clothing. Flush skin with water. If skin irritation occurs, get medical advice/attention. Wash contaminated clothing before reuse.

IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing. Call the Poison Centre or a doctor if you feel unwell. Apply artificial respiration if not breathing.

EMERGENCY RESPONSE

In a transport emergency call 111, Police or Fire Brigade. For specialist advice in a chemical emergency call 0800 CHEMCALL (0800 243 622).

SAFETY DATA SHEET

Additional information is provided in the safety data sheet. Phone 09 273 3456 or visit www.agpro.co.nz for a copy.

PROTECTION OF ANIMALS

Store securely away from animals as accidental poisoning may be fatal. Keep poultry, pets, dogs and stock off treated areas. **AGPRO MOLLUS** contains a deterrent which should stop non-target animals from consuming it. There may be some animals that are not deterred. If accidental poisoning occurs, consult a veterinarian.

GENERAL INFORMATION

AGPRO MOLLUS controls slugs and snails by irreversibly destroying cells that produce mucus which is essential for movement. Optimum results are obtained when applied under mild damp conditions when slugs and snails are most active. Avoid/delay application if rain is imminent. A repeat application may be necessary where infestation is high.



RESISTANCE MANAGEMENT

Resistance to metaldehyde may occur with repeated use. There are no resistance management strategies for metaldehyde. Use in accordance with label directions.

WITHHOLDING PERIOD

Do not apply to plant parts either for human consumption or those that will be grazed by or fed to stock. Withhold stock for 7 days following application.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

DIRECTIONS FOR USE

TIMING

Apply baits evenly using calibrated ground application equipment. Application should be targeted when crops are most vulnerable. In broadacre crops this is usually from drilling to emergence. For other crops, time the application to protect sensitive growth stages.

- Do not apply to plant parts either for human consumption or those that will be grazed by or . fed to stock.
- Do not place in heaps.
- Use the higher rates where pest pressure is moderate or higher or under heavy rainfall conditions.

APPLICATION RATE

The maximum application rate is 8kg/ha (0.4kg a.i./ha) except for when the substance is used on container plants whereby no maximum application rate is set.

For the control of Slugs and Snails.			
CROP	APPLICATION RATE	CRITICAL COMMENTS	25/09/2023
Cereals, fodder crops, other field crops		Broadcast evenly using a granule/bait spreader over soil surface or within a band of the area to be treated. Apply when crops are most vulnerable — most likely from drilling to emergence.	_,_,_
Vegetable crops	Broadcast Application: 4 - 8 kg/ha	Broadcast evenly over the soil surface of the area. Do not apply to edible plant parts. For short rotation crops such as Lettuce, apply pre-sowing or transplant during periods when crops are likely to be affected by pests (early spring to late autumn and during warm winters). Optimum results are obtained when applying before crop is present and also minimises potential for baits to lodge in edible foliage.	
Newly renovated pastures		Broadcast evenly using a granule/bait spreader over soil surface or within a band of the area to be treated. Apply when crops are most vulnerable — most likely from drilling to emergence.	

CONDITIONS OF SUPPLY

As the use of this product is beyond the control of AGPRO NZ Ltd, no warranty expressed or implied, beyond that covered under the Fair Trading Act is given by AGPRO NZ Ltd regarding its suitability, fitness, or efficacy for any purpose for which it is used by the user, whether in accordance with the directions or not. AGPRO NZ Ltd accepts no responsibility for any consequence whatsoever resulting from the use of this product.

Batch Number: Date of Manufacture:

Artwork Reference Number

