FLAMMABLE LIQUID DANGER ECOTOXIC Keep Out of Reach of Children

NB: Read the directions and information on the label before using

Glacier

Contains 250 g/L difenoconazole in the form of an emulsifiable concentrate. Also contains 30 g/l N-methyl-2-pyrrolidone and 610 g/l Xylene.

Glacier is a fungicide for the control of Stemphylium leaf spot in Asparagus, Leaf spots in Carrots, Early blight in Potatoes and Ring spot Vegetable brassicas.

Registered pursuant to the ACVM Act 1997, No. P8103 See <u>www.foodsafety.govt.nz</u> for registration conditions.

Approved pursuant to HSNO Act 1996, No. HSR100141 See <u>www.epa.govt.nz</u> for approval conditions.

Pack Size: 20 litres





WARNING:

- Warning: Flammable liquid and vapour.
- Warning: May be harmful if swallowed, inhaled or absorbed through skin.
- Warning: May cause skin irritation.
- Warning: May cause eye irritation.
- Danger: May damage fertility or the unborn child.
- Warning: May cause organ damage from repeated oral exposure at high doses.

Warning: Very toxic to aquatic life.

Approved Handler: This product must be under the control of an approved handler during use.

Record Keeping: Records must be kept for 3 years.

PRECAUTIONS:

Personal Protection

Do not eat, drink or smoke whilst using.

Wear impermeable gloves, coat, hat and trousers during use.

Immediately after use or before any meal break, remove protective clothing and thoroughly wash hands and face.

May be harmful if swallowed. Avoid skin and eye contact.

Storage

Store in original container tightly closed and in a locked dry, cool, well ventilated area away from foodstuffs. Under appropriate storage, the product shows no significant degradation within 2 years from the date of manufacture. Contact the supplier if product is older than 2 years.

Spills

Wear the above mentioned protective clothing when dealing with spills. Contain the spill and recover spilt product with sawdust or other inert absorbent material then place recovered material into properly labelled drum. Deal with spillages immediately and prevent contamination of watercourses, drains, streams, ponds. If this is likely to occur, warn the local authorities immediately.

Disposal

Dispose of product by following label directions or at an approved landfill. Avoid contamination of waterways with product or empty container.

Container Disposal

Triple rinse container and add rinsate to spray tank. Burn rinsed container if circumstances, especially wind direction, are suitable. Otherwise bury in suitable landfill.



FIRST AID:

Contact National Poisons and Hazardous Chemicals Information Centre ph 0800 764 766 (0800 POISON) or a doctor for medical advice.

If swallowed do not induce vomiting. Seek medical advice immediately.

If skin contact occurs, remove contaminated clothing and flush skin with soap and running water immediately. Wash contaminated clothing before re-use.

If eye contact occurs, wash out eyes with running water immediately.

GENERAL INFORMATION:

Glacier is a foliar applied systemic fungicide from the DMI group with eradicant and protective activity against a range of foliar diseases in Asparagus, Carrots, Potatoes and Vegetable brassicas.

DIRECTIONS FOR USE:

It is an offence to use this product on animals.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Withholding Period:

Asparagus:	Nil
Carrots:	14 days
Potatoes:	14 days
Vegetable brassicas:	14 days

Application:

Ground Application. Apply using conventional spray equipment in sufficient water to obtain complete coverage without run-off.

Mixing: Add the required quantity of Glacier to a half filled spray tank the complete filling. Maintain agitation throughout filling and application.

Rainfall: Do not apply if rain is expected within 3 hours of application.

Compatibility: Glacier is compatible with most commonly used fungicides and insecticides.

Resistance: Resistance to Glacier and other related DMI fungicides could develop from repeated use. Use strictly according to label directions to minimise the risk of resistance developing. Apply a maximum of **three** DMI fungicides per crop per season at full label rates.

Crop	Target	Rate	Remarks
Asparagus	Stemphylium leaf spot	500ml/ha	After close up of beds, apply at the first sign of the disease and repeat at 3 week intervals with a maximum of 3 applications during active fern growth. The addition of a non-ionic wetting agent at manufacturer's label recommendations is essential to assist coverage. Apply an approved protectant fungicide at other times. Aerial application - apply in a minimum of 100 litres water/ha to achieve satisfactory coverage.
Carrots	Leaf Spots	in	Apply at the first sign of the diseases and repeat at 2 to 3 week intervals with a
	(Alternaria & Cercospora)	500 - 700 l water/ha	maximum of 3 applications during active crop growth. Apply an approved protectant fungicide at other times.
Potatoes	Early blight		Apply at the first signs of the disease and depending on disease pressure, repeat at 2 to 3 week intervals with a maximum of 3 applications during active crop growth. Apply an approved protectant fungicide at other times.
Vegetable brassicas	Ringspot		Apply at the first sign of disease and repeat at 2 week intervals with a maximum of 3 applications. Apply an approved protectant fungicide at other times.



CONDITIONS OF SALE:

The use of Glacier being beyond the control of the manufacturer, no warranty expressed or implied is given by AGPRO NZ LTD regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and AGPRO NZ LTD accepts no responsibility for any consequence whatsoever resulting from the use of this product.

Date of Manufacture: Batch Number:

Registered to and distributed by: AGPRO NZ LTD PO BOX 58-963, GREENMOUNT 10 POLARIS PLACE, EAST TAMAKI AUCKLAND 2163 PH 0-9-273 3456 FAX 0-9-273 3457 Web: www.agpro.co.nz

