

**KEEP OUT OF REACH OF CHILDREN**

Read label before use

# AGPRO® Glyphosate Gel

**Contains 120 g/kg glyphosate isopropylamine salt in the form of a gel**

A ready to use herbicide gel for the direct application onto cut stems



**5L**

Approved pursuant to HSNO Act 1996,  
Approval Code HSR100736  
See [www.epa.govt.nz](http://www.epa.govt.nz) for approval controls

Manufactured and distributed by:

**AGPRO (N.Z.) Limited**

10 Polaris Place, East Tamaki, Auckland 2013

PO Box 58-963, Botany, Auckland 2163

Ph 09 273 3456

[www.agpro.co.nz](http://www.agpro.co.nz)

**AGPRO**

LBL 111.1

## HAZARD CLASSIFICATION

Designed for biocidal action

**This product must not be used for any purpose, or in any manner, contrary to this label unless authorised under appropriate legislation.**

## ECOTOXICITY

May cause long lasting harmful effects to aquatic life  
Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area. Avoid contamination of any water supply with product or empty container.

## PRECAUTIONS

Non-Selective Herbicide – very toxic to some plant species.

## PERSONAL PROTECTION

Avoid contact with skin and eyes or ingestion. When applying this product chemically resistant gloves, overalls, boots and eye protection should be worn. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work. Wash all protective clothing after work.

## STORAGE

Store in original container, tightly closed, in a secure, dry, cool and well-ventilated area avoiding direct sunlight and away from foodstuffs, fertilisers or seeds. Storage must be in accordance with the current version of the NZ Standard for the Management of Agrichemicals (NZS 8409).

## SHELF LIFE

When stored appropriately, this product should show no significant degradation for 24 months from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

## DISPOSAL

**PRODUCT DISPOSAL:** Dispose of this product only by using according to the label directions or if this is not possible contact your supplier for advice.

**CONTAINER DISPOSAL:** Recycle clean containers at an Agrecovery depot, or puncture and dispose of at an approved landfill. Do not burn. Do not use container for any other purpose.

Batch Number:

Date of Manufacture:

## FIRST AID

For advice call the National Poisons Centre 0800 POISON (0800 764 766) (24 hours) or a doctor. If medical advice is needed have product container or label to hand.

**IF SWALLOWED:** Do NOT induce vomiting Call the Poisons Centre or a doctor if you feel unwell.

**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If irritation persists, get medical advice/attention.

**IF ON SKIN OR HAIR:** Remove contaminated clothing. Flush skin with water. If skin irritation occurs, get medical advice/attention. Wash contaminated clothing before reuse.

**IF INHALED:** Remove to fresh air and keep at rest in a position comfortable for breathing. Call the Poison Centre or a doctor if you feel unwell. Apply artificial respiration if not breathing.

## SPILLAGE

Exclude non-essential people from the area. Wear protective clothing and prevent chemical from entering waterways. Scrape up and place in a sealable and labelled waste container. Wash area and absorb with further inert material. Dispose of safely at an approved landfill.

## SAFETY DATA SHEET

Additional information is provided in the safety data sheet. Phone 09 273 3456 or visit [www.agpro.co.nz](http://www.agpro.co.nz) for a copy.

## GENERAL INSTRUCTIONS

**AGPRO Glyphosate Gel** is a ready to use herbicide for direct application onto a wide range of trees, woody weeds, shrubs, scrambling perennial weeds and weeds with creeping or bulbous roots. The product is applied by swabbing the cut stump or stem using the specially designed applicator. Depending on the size of the plant there is usually no re-sprouting after application and the plant will die over the following weeks. On large trees this may take some months. Repeat application may be required if regrowth occurs.

**AGPRO Glyphosate Gel** may be applied at any time of year, however the best results will occur when the plant is actively growing, (spring to autumn). Rainfall during or within 12 hours after application will give a poor result.

## RESISTANCE MANAGEMENT

GROUP	9	HERBICIDE
-------	---	-----------

**AGPRO Glyphosate Gel** contains glyphosate as the isopropylamine salt. Glyphosate is a glycine herbicide which works by inhibiting EPSP synthase. Naturally occurring

resistant biotypes are known to exist and can become more prevalent after many years of continuous or excessive use of this product. To prevent or delay resistance use **AGPRO Glyphosate Gel** in rotation with herbicides that have a different mode of action. Refer to [www.resistance.nzpps.org](http://www.resistance.nzpps.org) for further resistance management advice.

## DIRECTIONS FOR USE

**Woody weeds:** These include old man's beard, climbing Spindleberry, Japanese honeysuckle, grey Willow, gorse, cotoneaster, Darwin's barberry, Elaeagnus, Woolly nightshade and Wilding Pines. Cut stems horizontally at 20mm and preferably no higher than 100mm above ground level. Apply a 3-5mm thick layer of **AGPRO Glyphosate Gel** over the cut surface remaining on the plant. On stems of 20mm diameter or greater, apply at least a 5mm thick layer in the case multi-stem plants treat 80 percent of the stems including all main stems.

**Woolly nightshade trees:** Cut a series of blazes 15-20mm deep around main trunks of the tree using an axe or pruning saw. Blazes should be evenly spaced with a 20-40mm gap between the blazes. Apply a 5mm layer of **AGPRO Glyphosate Gel** over the lower cut surface of the blaze.

**Rhizomatous plants:** These include Kahili ginger and agapanthus. On agapanthus prune all shoots horizontally at the rhizome and apply a 3-5mm layer of **AGPRO Glyphosate Gel** across the cut surface on the rhizome. On Kahili ginger prune and treat the same as the agapanthus, but if there are more than four rhizomes between the shoots, also drill a 10mm hole into every fourth rhizome from a shoot to 80 percent of the depth of the rhizome, and fill the hole with **AGPRO Glyphosate Gel**.

**Wandering Willie:** Apply **AGPRO Glyphosate Gel** at 16g per square metre over the foliage using a paint roller or similar making sure the foliage has been completely flattened during application.

## CONDITIONS OF SALE

As the use of this product is beyond the control of AGPRO (N.Z.) Limited, no warranty expressed or implied, beyond that covered under the Fair Trading Act is given by AGPRO (N.Z.) Limited regarding its suitability, fitness, or efficacy for any purpose for which it is used by the user, whether in accordance with the directions or not. AGPRO (N.Z.) Limited accepts no responsibility for any consequence whatsoever resulting from the use of this product.



AGPRO is a registered trademark of AGPRO (N.Z.) Limited.

LBL 111.1