

**WARNING**  
**KEEP OUT OF REACH OF CHILDREN**  
**TOXIC**  
**ECOTOXIC**

**SABLE**

GROUP	A	HERBICIDE
-------	---	-----------

Contains 69 g/L fenoxaprop-P-ethyl in the form of an oil in water emulsion

For selective grass weed control in autumn and spring sown Wheat, Perennial Ryegrass and Pasture

This product must be under control of an Approved Handler when wide dispersive use, or used by a professional contractor.

**READ LABEL BEFORE USE**



Registered pursuant to the ACVM Act 1997 No: P009478  
See [www.nzfa.govt/acvm](http://www.nzfa.govt/acvm) for registration conditions  
Approved pursuant to HSNO Act 1996 No HSR101234  
See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions

Date of manufacture:  
Batch no:  
Net contents: 10L

**Distributed by:**  
AGPRO NZ Ltd  
10 Polaris Place  
East Tamaki  
P O Box 58-963  
East Tamaki  
Auckland 2163  
Ph 09 273 3456  
Fax 09 273 3457  
Website: [www.agpro.co.nz](http://www.agpro.co.nz)

**HSNO CLASSIFICATION**

6.5B, 6.9B, 9.1B

**WARNING**

May cause an allergic skin reaction.

May cause damage to organs through prolonged or repeated oral exposure.

Toxic to aquatic life with long lasting effects.

**PRECAUTIONS**

Do not breathe vapors, mist or spray. Avoid release to aquatic environment. Apply product carefully. Spray drift may cause serious damage to desirable plants.

**PERSONAL PROTECTION:** Avoid skin or eye contact or inhalation. Do not eat, drink or smoke while using. Wear protective clothing such as waterproof hat, coat and trousers when using. Wear eye protection and gloves. Remove protective clothing and wash hands and face thoroughly before meals and after work. Contaminated work clothing should not be allowed out of the workplace. Wash contaminated clothing before reuse.

**STORAGE:** Store in original container tightly closed and in a secure dry, cool, well-ventilated area, away from foodstuffs, seeds and fertilizers.

**SPILLS AND DISPOSAL:** Wear personal protective clothing and equipment as described in the PERSONAL PROTECTION section. Contain the spill then recover spilt product by absorbing with sawdust or an inert absorbent material. Transfer the recovered material to a labeled drum that can be sealed.

Deal with all spillages immediately. If contamination of drains, streams, watercourses is unavoidable, advise emergency services and Regional Authority.

**DISPOSAL:** Dispose of this product only by using according to the label or to an approved landfill. Avoid contamination of any water supply with product or empty container.

**CONTAINER DISPOSAL:** Triple rinse container and add washings to spray tank. Burn if circumstances, especially wind direction, permit. Otherwise bury in a suitable land fill.

**FIRST AID**

If medical advice is needed, have product container or label at hand. For advice, call the National Poisons Centre 0800 POISON (0800 764766) or a doctor.

If skin contact occurs, wash with plenty of soap and water. If skin irritation or rash occurs get medical advice.

**SAFETY DATA SHEET**

If additional hazard information is required, refer to the Safety Data Sheet. Copy available from supplier.

**SHELF LIFE**

When stored appropriately, this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

**WEEDS CONTROLLED**

Wild oats, lesser canary grass (*Phalaris minor*), yellow bristle grass (*Setaria glauca*).

**RESISTANT WEEDS**

Gnawed canary grass (*Phalaris paradoxa*), annual and hybrid ryegrasses, tall fescue, chewings fescue, cocksfoot, browntop, grazing brome (spiky broome; *Bromus stamineus*), prairie grass and most broadleaved weeds

**RESISTANT WEED WARNING**

SABLE is a **GROUP A HERBICIDE** from the aryloxyphenoxypropionate (FOPs) chemical group. Naturally occurring weed biotypes resistant to aryloxyphenoxypropionate herbicides can become more prevalent after many years of continuous use. To delay the onset of, or control resistant weeds, use in rotation with herbicides of a different mode of action. Since the occurrence of resistant weeds is difficult to detect prior to

use, Shandong Rainbow International Co. Ltd accepts no liability for losses that may result from the failure of this product to control resistant weeds.

**MIXING**

**Invert container several times and** shake container well before use. Add the required amount of SABLE to a partly filled spray tank with agitation system operating. Complete filling the tank with water and apply. After spraying, flush the spray system with clean water.

**APPLICATION**

For application by ground-based application methods only. Do not apply onto or into water. Apply at no more than maximum rate of 53 g fenoxaprop-P-ethyl /hectare, once a year. Do not overlap swath width in ryegrass crops as yield suppression may occur. Flush spray system with clean water after use. Reentry period is 24 hours. Do not enter treated area within this time unless wearing personal protective equipment.

**DIRECTIONS FOR USE**

Crop	Weeds	Rate	Remarks
<b>Wheat, Barley</b>	Wild oats	750 mL/hectare in 250 litres of water. Add adjuvant oil at recommended rates	Apply when wild oats are between 3 leaf and 1 <sup>st</sup> node visible growth stage and when they are growing actively.
	Lesser canary grass ( <i>Phalaris</i> )		Apply when weeds are between the 3 leaf visible stage and the end of tillering and are growing actively. Avoid application during periods of moisture stress.
<b>Perennial ryegrass</b> ( <i>Lolium perenne</i> )	Wild oats	750 mL/hectare in 250 litres of water. <b>DO NOT use with any other agrochemicals or adjuvants.</b>	Apply in early spring when wild oats are between 3 leaf and end of tillering growth stage and growing actively. Note: Seed yield response is likely to be reduced the longer the delay in the removal of wild oats in spring.
<b>Pasture</b>	Yellow bristle grass ( <i>Setaria glauca</i> )	750 mL/hectare in 250 litres of water. <b>DO NOT use with any other agrochemicals or adjuvants.</b>	Apply to established pasture 7 days after grazing during the period of December/January.

**RAINFASTNESS**

SABLE is rainfast within one hour of application.

**SOIL ACTIVITY**

SABLE is rapidly broken down by soil microbes causing no restrictions on crop rotations.

**UNDERSOWN CROPS**

SABLE can be used on crops undersown with certain grasses (Refer Resistant Weeds section) and clover. Do not use on crops undersown with "Concord" ryegrass.

**COMPATIBILITY**

DO NOT USE SABLE IN ANY MIXTURES WHEN APPLYING TO PERENNIAL RYEGRASS. When used on wheat, SABLE can be tank mixed with approved expoxiconazole and chlormequat-chloride products at label rates in two and three way mixes. Care must be taken to ensure that recommended timings for all products are followed. The addition of a suitable adjuvant oil at label rates is recommended. DO NOT use SABLE within 5 days of broad-leaved herbicide applications.



**WITHHOLDING PERIODS**

**Wheat** -70 days (Do not graze treated crop for at least 70 days after treatment)

**Perennial ryegrass** -42 days (Do not graze treated crop for at least 42 days after treatment)

**Pasture** – 28 days (Do not graze treated pasture for at least 28 days after treatment)

**It is an offence for users of this product to cause residues exceeding the relevant MRL in Food**  
**Notice: Maximum Residue Levels for Agricultural Compounds.**

**Registered to:**

Shandong Rainbow International Co., Ltd  
19<sup>th</sup> & 20<sup>th</sup> Floor, Hanyu Financial Centre,  
Building A3-4, No. 7000 East Jingshi Road,  
Jinan, China 250101

**NZ Agent:**

AR & JA DRYSDALE LTD  
PO Box 72275  
Papakura 2244  
Tel: (021) 939 728

**CONDITIONS OF SALE**

The use of SABLE herbicide being beyond the control of the suppliers, no warranty expressed or implied, beyond that covered under the Fair Trading Act is given by the supplier regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and the supplier accepts no responsibility for any consequences whatsoever resulting from the use of this product.