

HARMFUL
Keep out of reach of children
ECOTOXIC

TRIPLUS™
Fungicide

Contains 500g/kg trifloxystrobin in the form of a water dispersible granule

For the control of black spot and powdery mildew in apples, black spot in pears, powdery mildew in winter squash and pumpkins, Sclerotinia in kiwifruit, brown rot in stonefruit (except cherries) and scab in citrus

Note: Please read all the label information before using this product.

Registered pursuant to the ACVM Act 1997 No: P009147
See www.nzfa.govt/acvm for registration conditions
Approved pursuant to HSNO Act 1996 No HSR000642
See www.epa.govt.nz for approval conditions

Date of manufacture:
Batch no:
Net contents: 1 kg

Distributed by:
AGPRO NZ Ltd
10 Polaris Place
East Tamaki
P O Box 58-963
East Tamaki
Auckland 2163
Ph 09 273 3456
Fax 09 273 3457
Website: www.agpro.co.nz

HSNO CLASSIFICATION

6.5B, 6.9B, 9.1A

WARNING

Harmful - may cause sensitisation from prolonged skin contact.

Harmful - presumed to/may cause organ damage from repeated oral exposure at high doses.

Toxic to aquatic organisms - must not be applied directly onto or into water.

PRECAUTIONS

PERSONAL PROTECTION: Do not eat, drink or smoke while using. Wear protective clothing such as impervious gloves, waterproof hat, coat and trousers when using. Remove protective clothing and wash hands and face thoroughly before meals and after work. May be harmful if swallowed, inhaled or absorbed through the skin. Avoid skin contact or inhalation.

STORAGE: Store in original container tightly closed and in a locked dry, cool, well-ventilated area, away from foodstuffs. When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of this product if it is older than this.

SPILLS: When dealing with spills wear personal protective clothing and equipment as described in the section. Sweep up spilled material, collect and store in properly labeled, sealed drums for safe disposal. Avoid creating dust. If spray mixture is spilled, contain and absorb in suitable inert absorbent material and collect in drums as above. Deal with all spillages immediately. If contamination of drains, streams, watercourses, etc. is unavoidable, warn the local water authority.

DISPOSAL: Dispose of this product only by using according to the label or at an approved landfill. Avoid contamination of any water supply with product or empty container. Ensure container is completely empty. Burn if circumstances, especially wind direction, permit. Otherwise bury in a suitable land fill.

FIRST AID

If swallowed do not induce vomiting. If splashed in eyes, wash out immediately with running water for several minutes. If skin or hair contact occurs, remove contaminated clothing and wash with soap and running water. If inhaled, immediately move to fresh air and rest. Maintain half upright position if breathing is difficult. Call the National Poisons Centre 0800 POISON (0800 764 766) (24hrs) or a doctor immediately for advice.

EMERGENCY RESPONSE

In a transport emergency call 111, Police or Fire Brigade. For specialist advice in an emergency only, call 0800 CHEMCALL (0800 243 6225) (24hrs).

GENERAL INFORMATION

TRIPLUS is a mesostemic strobilurin fungicide with preventative properties for the control of black spot and powdery mildew in apples, black spot in pears, powdery mildew in winter squash and pumpkins, Sclerotinia in kiwifruit, brown rot in stonefruit (except cherries) and scab in citrus.

TRIPLUS is a synthetic analogue of a naturally occurring fungicide produced by a wood decaying fungi.

TRIPLUS is compatible with IMC programmes based on *Typhlodromus pyri*.

RESISTANCE

Resistance to this fungicide and related strobilurin in products could develop in some disease situations from repeated use. To minimise this risk, use strictly in accordance with the label instructions, and in pipfruit apply no more than two sequential applications of two sprays with a maximum of four applications per season. In winter squash, pumpkins and citrus apply no more than two applications of TRIPLUS or other strobilurin fungicides per season. Follow individual crop resistance management strategies.

DIRECTIONS FOR USE

The rates suggested are for high volume spraying of a dilute spray without excessive run-off. For concentrate spraying adjust dilution rate accordingly. Do not add wetting agents to TRIPLUS.

Crop	Disease	Rate / Remarks
Apples	Black Spot, Powdery mildew	Apply 10g/100 litres (17.5ml measure) of water. DO NOT add other products (including wetting agents) when used over the flowering - early fruit set period.
Pears	Black spot	Apply as for apples.

TIMING

For apples and pears apply TRIPLUS as a preventative application at 7-9 day intervals during the period covering flowering to the end of extension growth.

The 7 day interval is recommended when conditions are conducive to severe Black Spot infection and during periods of rapid tree growth. If the interval exceeds 9 days and there has been a recent infection period, the addition of a DMI black spot fungicide product is required, used according to label recommendations. If a protectant fungicide follows a TRIPLUS application, it should be applied no later than 7 days after the last TRIPLUS spray.

For Resistance Management apply TRIPLUS in a maximum of two consecutive applications but do not exceed four total applications per season.

Crop	Disease	Rate / Remarks
Winter squash, Pumpkins	Powdery mildew	Apply 125g/ha (220ml measure) in sufficient water to obtain uniform coverage. Apply no more than 2 applications, 14 days apart.
Kiwifruit	Sclerotinia	Apply 15g/100 litres (26.25ml measure) of water (minimum 300g/ha) during flowering.
Stonefruit (except cherries)	Brown rot	Apply 12.5g/100 litres(22ml measure) (minimum 250g/ha) in a regular blossom programme from pink to petal fall inclusive. Thereafter apply alternative non-QoI fungicides through until harvest. For resistance management apply no more than 3 applications per season.
Citrus	Scab	Apply 10g/100 litres (17.5ml measure) as a full cover spray to the point of run-off. Make 2 applications, the first application at flowering/fruit set and the second up to 1 month after fruit set.

It is an offence to use this product on animals.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds

MIXING

TRIPLUS mixes readily with water. Add the required amount of TRIPLUS to the spray tank filled to one-sixth of its capacity. Under agitation add the remainder of the water and ensure complete mixing before spraying. Continue agitation during spraying and after a stoppage before recommencing spraying.

COMPATIBILITY

TRIPLUS is compatible with most of the commonly used insecticides and fungicides. DO NOT add other products to TRIPLUS when used over the flowering - early fruit set period.

WITHHOLDING PERIOD

Apples, pears	70 days
Winter squash, pumpkins	14 days
Kiwifruit	Do NOT apply after flowering
Stonefruit (except cherries)	Do NOT apply after petal fall
Citrus (multi flowering)	28 days
Citrus (Clementine/Satsuma mandarins)	120 days

Registered to:

GLOBAL AGRICHEM LIMITED
Suite 4, Level 1
38 Whitaker Place
Grafton
Auckland
Phone 022 191-5641

CONDITIONS OF SALE

The use of TRIPLUS being beyond the control of the manufacturer, no warranty expressed or implied is given by manufacturer/distributor regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not, and the manufacturer/distributor accepts no responsibility for any consequence whatsoever resulting from the use of this product.